

NOTE

ON LABELED VERTEX-TRANSITIVE DIGRAPHS WITH A PRIME NUMBER OF VERTICES

Chong-Yun CHAO

University of Pittsburgh, Pittsburgh, PA 15260, USA

Jacqueline G. WELLS

Pennsylvania State University, McKeesport, PA 15132, USA

Received 23 July 1982

Revised 26 November 1982

We enumerate, up to isomorphism, several classes of labeled vertex-transitive digraphs with a prime number of vertices.

There are many unsolved enumeration problems stated in [5]. Recently, Robinson in [8] posed more enumeration problems. Here, we give some partial answers to the problems posed on p. 181 in [8], i.e., by using some of the results in [3] and [4], we enumerate several classes of labeled vertex-transitive digraphs with a prime number, p , of vertices; specifically, we count those that are symmetric (both vertex-transitive and edge-transitive), those with a given group of automorphisms and those that are self-complementary. Roughly, our method is to use the group of automorphisms, Pólya's and de Bruijn's enumeration theorems and the following lemma.

Lemma 1 (See [5, Chapter 1]). *Let H_1, H_2, \dots, H_t be all nonisomorphic digraphs (graphs) with n vertices and $|\Gamma(H_i)|$ be the order of the automorphism group $\Gamma(H_i)$ of H_i for $i = 1, 2, \dots, t$. Then the total number of distinct labeled digraphs (graphs) with n vertices is $\sum_{i=1}^t n!/|\Gamma(H_i)|$.*

Theorem 1. *The number of labeled symmetric digraphs on a prime number p vertices is*

$$2 + \sum_n \frac{(p-1)!}{n}$$

where n is a divisor of $p-1$, and $1 \leq n < p-1$.

Proof. In [3], Theorem 3 states that the group of automorphisms $\Gamma(X)$ of a non-null and non-complete vertex-transitive digraph X on p vertices is generated by R and σ with defining relations $R^p = e$, $\sigma^n = e$ and $\sigma^{-1} R \sigma = R^r$ where e is the identity of $\Gamma(X)$, n divides $p-1$, $1 \leq n < p-1$ and $r^n \equiv 1 \pmod{p}$. If the digraph

X is symmetric, then, by Corollary 3.1 in [3], $|\Gamma(X)| = np$ where n is the degree of X .

By Lemma 1, including the null and complete graphs with p vertices, the total number of labeled symmetric digraphs with p vertices is

$$2 + \sum_n \frac{p!}{np} = 2 + \sum_n \frac{(p-1)}{n} \quad (1)$$

where n is a divisor of $p-1$, and $1 \leq n < p-1$.

In Theorem 4 of [4] we used the group structure described in the preceding proof to enumerate the non-isomorphic digraphs with a prime number p of vertices and with group of automorphisms $\cong \langle R, \sigma \rangle$, where $\sigma = \tau^m$ and $\tau^{p-1} = e$. This enumerating function is

$$\frac{1}{m} \sum_d \phi(d)(1 + x^{dnp})^{m/d} \quad (1)$$

where the sum is taken over the divisors d of m , $mn = p-1$ and where ϕ is Euler's ϕ function.

Lemma 2. *The enumerating function for the non-isomorphic classes of vertex-transitive digraphs with a prime number of vertices and with the group of automorphisms $\langle R, \tau^m \rangle$ is*

$$\sum_D \mu\left(\frac{m}{D}\right) \cdot \frac{1}{D} \sum_d \phi(d)(1 + x^{dnp})^{D/d} \quad (2)$$

where m divides $p-1$, $Dn = (p-1)$, the first summation is taken over the divisors D of m and the second over the divisors d of D , μ is the Möbius function.

We omit the proof here. Our Lemma 2 is a generalization of Theorem 5 in [4]. There we enumerated the digraphs with p vertices and with the group of automorphisms $\langle R, \tau^{p-1} \rangle = \langle R \rangle$. Here we apply the same exclusion-inclusion principle to the group of automorphisms $\langle R, \tau^m \rangle$.

Example 1. For $p = 13$ and $m = 6$, we enumerate the digraphs with 13 vertices each of whose group of automorphisms is $\langle R, \tau^6 \rangle$. By (2), we have

$$\begin{aligned} & \frac{1}{6} \sum_{d|6} \phi(d)(1 + x^{26d})^{6/d} - \frac{1}{3} \sum_{d|3} \phi(d)(1 + x^{52d})^{3/d} \\ & - \frac{1}{2} \sum_{d|2} \phi(d)(1 + x^{78d})^{2/d} + (1 + x^{156}) = x^{26} + 2x^{52} + 3x^{78} + 2x^{104} + x^{130}. \end{aligned}$$

Thus, on 13 vertices there are 9 nonisomorphic digraphs with the group of automorphisms $\langle R, \tau^6 \rangle$: one has 26 edges, 2 have 52 edges, etc. Since the order of the group $\langle R, \tau^6 \rangle$ is $2 \cdot 13$, by Lemma 1, the number of labeled digraphs on 13 vertices having the group $\langle R, \tau^6 \rangle$ is $9 \cdot 13!/2 \cdot 13 = 54 \cdot 11!$.

Replacing x by 1 in (2) and applying Lemma 1, we have

Theorem 2. *The number of labeled digraphs with a prime p vertices and with the group of automorphisms $\langle R, \tau^m \rangle$ is*

$$\left[\sum_D \mu\left(\frac{m}{D}\right) \cdot \frac{1}{D} \sum_d \phi(d) \cdot 2^{D/d} \right] m(p-2)! \quad (3)$$

where m divides $p-1$, the first summation is taken over the divisor D of m and the second summation is taken over the divisor d of D .

A vertex-transitive digraph is said to be strongly vertex-transitive if its group of automorphisms is cyclic.

Corollary 2.1. *The number of labeled strongly vertex-transitive digraphs with a prime number p vertices is*

$$\left[\sum_{D|(p-1)} \mu\left(\frac{p-1}{D}\right) \frac{1}{D} \sum_{d|D} \phi(d) \cdot 2^{D/d} \right] (p-1)! \quad (4)$$

Proof. It follows from (3) with $m = p-1$.

Finally, we shall count the labeled self-complementary digraphs with p vertices each of whose group is $\langle R, \tau^m \rangle$. We showed in [3] that a digraph with p vertices and with the group of automorphisms $\langle R, \tau^m \rangle$ has degree a multiple of $(p-1)/m$, since τ^m fixes one vertex and is a collection of cycles of length $(p-1)/m$. A self-complementary vertex-transitive digraph with p vertices has degree $\frac{1}{2}(p-1)$. Thus, $\frac{1}{2}(p-1)$ is a multiple of $(p-1)/m$ and m must be even.

Theorem 3. *The number of labeled vertex-transitive self-complementary digraphs with a prime number p vertices and with the group of automorphisms $\langle R, \tau^m \rangle$ is*

$$\left[\sum_D \frac{\mu(m/d)}{D} \sum_d \phi(d) \cdot 2^{D/d} \right] m(p-2)! \quad (5)$$

where m is an even divisor of $p-1$, the first summation is taken over the even divisors D of m and the second summation is taken over the even divisors d of D .

Proof. We apply de Bruijn's Theorem [1] to the enumeration (2) of digraphs each of whose group is $\langle R, \tau^m \rangle$. By using a second permutation group which places each digraph in the same class with its complement as in [4] and [7], we are able to determine the number of those that are self-complementary. This number is

$$\sum_D \frac{\mu(m/D)}{D} \sum_d \phi(d) \frac{\partial^{D/d}}{\partial z_d^{D/d}} e^{2(z_2 + z_4 + \dots)}, \quad (6)$$

evaluated at $z_1 = z_2 = \dots = 0$. Since only even values of d result in non-zero partial derivatives, we combine (6) with Lemma 1 to obtain (5).

Corollary 3.1. *The number of labeled strongly vertex-transitive self-complementary digraphs with a prime number p vertices is*

$$\left[\sum_{D|p-1} \frac{\mu((p-1)/D)}{D} \sum_{d|D} \phi(d) \cdot 2^{D/d} \right] (p-1)! \quad (7)$$

where D and d are even.

Proof. It follows from (5) by letting $m = p - 1$.

Corollary 3.2. *The total number of labeled vertex-transitive self-complementary digraphs with a prime number p of vertices is*

$$\sum_{m|p-1} \left[\sum_{D|m} \frac{\mu(m/D)}{D} \sum_{d|D} \phi(d) \cdot 2^{D/d} \right] m(p-2)! \quad (8)$$

where m , D and d are even.

Proof. The total number is obtained by summing (5) over all even m .

Example 2. For $p = 13$ we count the self-complementary vertex-transitive digraphs with 13 vertices and with the group of automorphisms $\langle R, \tau^m \rangle$. For $m = 12$,

$$\begin{aligned} & \frac{1}{12} \sum_{d|12} \phi(d) \cdot 2^{12/d} - \frac{1}{6} \sum_{d|6} \phi(d) \cdot 2^{6/d} \\ & - \frac{1}{4} \sum_{d|4} \phi(d) \cdot 2^{4/d} + \frac{1}{2} \cdot 2 = 8 - 2 - 2 + 1 = 5. \end{aligned}$$

For $m = 6$,

$$\frac{1}{6} \sum_{d|6} \phi(d) \cdot 2^{6/d} - \frac{1}{2} \cdot 2 = 2 - 1 = 1.$$

For $m = 4$,

$$\frac{1}{4} \sum_{d|4} \phi(d) \cdot 2^{4/d} - \frac{1}{2} \cdot 2 = 2 - 1 = 1.$$

For $m = 2$,

$$\frac{1}{2} \cdot \phi(2) \cdot 2 = 1.$$

By (8), the total number of labeled self-complementary vertex-transitive digraphs on 13 vertices is

$$60 \cdot 11! + 6 \cdot 11! + 4 \cdot 11! + 2 \cdot 11! = 72 \cdot 11!$$

Finally, we observe that the total number of all labeled vertex-transitive digraphs with a prime number p vertices is obtained by summing (3) for all $m > 1$ such that m divides $p - 1$, and adding 2 to include the labeled null and complete graphs.

Note added in proof

A formula similar to our result in Lemma 2 was presented by Brian Alspach in "Point-symmetric graphs and digraphs of prime order and transitive permutation groups of prime degree", *J. Combin. Theory (B)* 15 (1973) 12–17.

References

- [1] N.G. de Bruijn, Pólya's theory of counting, in: E.F. Beckenbach, ed., *Applied Combinatorial Mathematics* (Wiley, New York, 1964) Ch. 5, 144–184.
- [2] C.Y. Chao, A note on labeled symmetric graphs, *Discrete Math.* 8 (1974) 295–297.
- [3] C.Y. Chao and J.G. Wells, A class of vertex-transitive digraphs, *J. Combin. Theory (B)* 14 (1973) 246–255.
- [4] C.Y. Chao and J.G. Wells, A class of vertex-transitive digraphs II, *J. Combin. Theory (B)* 32 (1982) 336–346.
- [5] F. Harary and E. Palmer, *Graphical Enumeration* (Academic Press, New York and London, 1973).
- [6] E.M. Palmer, The enumeration of graphs, in: L.W. Beineke and R.J. Wilson, ed., *Selected Topics in Graph Theory* (Academic Press, New York and London, 1978) Ch. 14, 385–416.
- [7] R.C. Read, Some applications of a theorem of de Bruijn, in: F. Harary, ed., *Graph Theory and Theoretical Physics* (Academic Press, New York and London, 1967).
- [8] R.W. Robinson, Counting graphs with a duality property, *London Math. Soc. Lecture Note Series*, Vol. 52 (Cambridge Univ. Press, Cambridge, 1981) 156–186.